# Clean Copy of the Claims

1. (currently amended) A compound of a formula below:

wherein

n is 0, 1, 2, or 3;

q is 0, 1, 2, or 3;

Y is a bond, C=O, or S(O); wherein t is 0, 1, or 2;

R<sup>1</sup> is selected from a group consisting of C<sub>1</sub>-C<sub>6</sub> alkyl, aryl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>1</sub>-C<sub>6</sub> alkylheterocyclic, C<sub>3</sub>-C<sub>8</sub> cycloalkyl, C<sub>1</sub>-C<sub>6</sub> alkylcycloalkyl, C<sub>1</sub>-C<sub>6</sub> alkylaryl, heterocyclyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, aryloxy, OC<sub>1</sub>-C<sub>6</sub> haloalkyl, -OC<sub>3</sub>-C<sub>8</sub> cycloalkyl, -OC<sub>1</sub>-C<sub>6</sub> alkylcycloalkyl, -NR<sup>7</sup>R<sup>8</sup>, -OC<sub>1</sub>-C<sub>6</sub> alkylaryl, -O-heterocyclic, and -OC<sub>1</sub>-C<sub>6</sub> alkylheterocyclic; and wherein each of cycloalkyl, aryl and heterocyclic group is optionally substituted with 1 to 3 groups independently selected from oxo, halo, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, C<sub>1</sub>-C<sub>6</sub> haloalkyl, CONR<sup>11</sup>R<sup>12</sup>, C<sub>0</sub>-C<sub>3</sub> alkylNR<sup>11</sup>R<sup>12</sup>, C<sub>0</sub>-C<sub>6</sub> alkylCOOR<sup>11</sup>, evano, and phenyl:

each  $R^{5}$  is selected from a group consisting of hydroxy, halogen,  $C_{1}$ - $C_{6}$  haloalkyl, aryl, heterocyclic, cyano,  $C_{1}$ - $C_{6}$  alkyl,  $C_{2}$ - $C_{6}$  alkonyl,  $C_{1}$ - $C_{6}$  alkoxy,  $-OC_{1}$ - $C_{6}$  haloalkyl,  $C_{0}$ - $C_{6}$  alkyl $NR^{7}R^{8}$ ,  $C_{0}$ - $C_{6}$  alkyl $COR^{7}$ ,  $C_{0}$ - $C_{6}$  alkyl $CO_{2}R^{7}$ ,  $NR^{7}SO_{2}R^{8}$ ,  $NR^{7}COR^{8}$ ,  $S(O)_{1}R^{7}$ , and  $-OC_{1}$ - $C_{6}$  alkylaryl wherein each of the aryl and heterocyclic groups is optionally substituted by oxo or alkyloxy;

R<sup>6</sup> is hydrogen or C<sub>1</sub>-C<sub>6</sub> alkyl;

each  $R^7$  is independently selected from a group consisting of hydrogen,  $C_1$ - $C_6$  alkyl,  $OC_1$ - $C_6$  alkyl,  $C_1$ - $C_6$  alkyl,  $C_1$ - $C_6$  alkyl,  $C_1$ - $C_6$  alkyl,  $C_1$ - $C_6$  alkyl, heterocyclic, and aryl, wherein each alkyl, is optionally substituted with 1-3 groups independently selected from  $C_1$ - $C_6$  alkoxy,  $SO_2R^{11}$ , and  $NR^{11}R^{12}$ ,

each  $R^8$  is independently selected from a group consisting of hydrogen,  $C_1\text{-}C_6$  alkyl, and aryl;

R9 is COR7 wherein R7 is as defined above:

R<sup>10</sup> is benzyl, optionally substituted with 1 or 2 groups selected from halo, C<sub>1</sub>-C<sub>6</sub>alkyl, haloalkyl, C<sub>1</sub>-C<sub>6</sub>alkoxy, and C<sub>1</sub>-C<sub>6</sub> haloalkoxyalkyl;

 $R^{11}$  and  $R^{12}$  are independently selected from a group consisting of hydrogen,  $C_1$ - $C_6$  alkyl, and arvl:

or a pharmaceutically acceptable salt thereof.

- 2. (previously presented) The compound according to Claim 1 wherein  $R^1$  is selected from a group consisting of  $C_1$ - $C_6$  alkoxy,  $C_1$ - $C_6$  alkylcycloalkyl,  $C_3$ - $C_8$  cycloalkyl,  $C_1$ - $C_6$  alkylheterocyclic, aryloxy, -OC<sub>1</sub>- $C_6$  haloalkyl, -OC<sub>3</sub>- $C_8$  cycloalkyl, -OC<sub>1</sub>- $C_6$  alkylaryl and -OC<sub>1</sub>- $C_6$  alkylheterocyclic wherein each of cycloalkyl, aryl and heterocyclic group is optionally substituted with 1 to 3 groups independently selected from oxo, halo,  $C_1$ - $C_6$  alkyl,  $C_1$ - $C_6$  alkoxy,  $C_1$ - $C_6$  haloalkyl, CONR<sup>11</sup>R<sup>12</sup> and  $C_0$ - $C_6$  alkylCOOR<sup>11</sup>.
- 3. (currently amended) A compound according to Claim I wherein  $R^1$  is selected from a group consisting of aryloxy,  $-OC_1-C_6$  haloalkyl,  $-OC_3-C_8$  cycloalkyl,  $-OC_1-C_6$  alkylaryl, Oheterocyclic, and  $-OC_1-C_6$  alkylheterocyclic; wherein each of cycloalkyl, aryl and heterocyclic group is optionally substituted with I to 3 groups independently selected from halo,  $C_1-C_6$  alkyl,  $C_1-C_6$  alkoxy,  $C_1-C_6$  haloalkyl, and  $C_0-C_6$  alkyl $COOR^{11}$ .
- 4. (previously presented) The compound according to Claim 1 wherein  $\mathbb{R}^1$  is selected from a group consisting of  $C_1\text{-}C_6$  alkyleycloalkyl,  $C_1\text{-}C_6$  alkylheterocyclic,  $C_3\text{-}C_8$  cycloalkyl and aryloxy, wherein each of cycloalkyl, aryl and heterocyclic group is optionally substituted with 1 to 3 groups independently selected from halo,  $C_1\text{-}C_6$  alkyl,  $C_1\text{-}C_6$  alkoxy,  $C_1\text{-}C_6$  haloalkyl, and  $C_0\text{-}C_6$  alkylCOOR<sup>11</sup>.
- 5. (currently amended) The compound according to Claim 1 Y is a bond; and  $R^1$  is alkylaryl, alkylheterocyclic,  $C_1$ - $C_6$  alkylcycloalkyl wherein the aryl, cycloalkyl and heterocyclic groups are each optionally substituted with 1, 2 or 3 groups independently selected from oxo, COOH,  $C_1$ - $C_6$  alkyl, and  $C_1$ - $C_6$  alkyoxy.

### 6-7. (canceled)

- 8. (currently amended) The compound of claim 1, wherein n is 0 or 1 and q is 1, 2, or 3.
- 9. (previously presented) The compound according to Claim 1 wherein n is 0 or 1; and  $\alpha$  is 2 or 3.

### 10-11. (canceled)

- (previously presented) A compound selected from the group consisting of:
  5-[Acetyl-(3,5-bis-trifluoromethyl-benzyl)-amino]-7-trifluoromethyl-2,3,4,5-tetrahydro-benzofblazenine-1-carboxylic acid isopropyl ester.
- 5-[(3,5-Bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-7-trifluoromethyl-2,3,4,5-tetrahydro-benzo[b]azepine-1-carboxylic acid isopropyl ester,
- 5-[(3,5-Bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-7-trifluoromethyl-2,3,4,5-tetrahydro-benzo[blazenine-1-carboxylic acid ethyl ester.
- 5-[Acetyl-(3,5-bis-trifluoromethyl-benzyl)-amino]-7-trifluoromethyl-2,3,4,5-tetrahydrobenzo[b]azepine-1-carboxylic acid ethyl ester.
- 5-[Acetyl-(3,5-bis-trifluoromethyl-benzyl)-amino]-2,3,4,5-tetrahydro-benzo[b]azepine-l-carboxylic acid isopropyl ester.
- 5-[(3,5-Bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2,3,4,5-tetrahydrobenzofblazepine-1-carboxylic acid isopropyl ester.
- 5-[(3,5-Bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-7-bromo-2,3,4,5-tetrahydro-benzo[b]azepine-1-carboxylic acid isopropyl ester,
- 5-[(3,5-Bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-7-bromo-2,3,4,5-tetrahydrobenzo[b]azepine-1-carboxylic acid ethyl ester,
- 5-[Acetyl-(3,5-bis-trifluoromethyl-benzyl)-amino]-7-bromo-2,3,4,5-tetrahydro-benzo[b]azepinel-carboxylic acid ethyl ester,
- 5-[Acetyl-(3,5-bis-trifluoromethyl-benzyl)-amino]-7-methoxy-2,3,4,5-tetrahydrobenzo[blazepine-1-carboxylic acid ethyl ester.
- 5-[Acetyl-(3,5-bis-trifluoromethyl-benzyl)-amino]-8-trifluoromethyl-2,3,4,5-tetrahydro-benzo[b]azepine-1-carboxylic acid isopropyl ester,
- 5-[Acetyl-(3,5-bis-trifluoromethyl-benzyl)-amino]-8-fluoro-7-trifluoromethyl-2,3,4,5-tetrahydrobenzo[b]azepine-1-carboxylic acid isopropyl ester.

- 4-[Acetyl-(3,5-bis-trifluoromethyl-benzyl)-amino]-7-trifluoromethyl-2,3,4,5-tetrahydro-benzo[b]azepine-1-carboxylic acid isopropyl ester,
- 5-[Acetyl-(3,5-bis-trifluoromethyl-benzyl)-amino]-8-chloro-2,3,4,5-tetrahydro-benzo[b]azepine-1-carboxylic acid isopropyl ester, and
- 5-[(3,5-Bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-8-chloro-2,3,4,5-tetrahydrobenzo[b]azepine-1-carboxylic acid isopropyl ester, or a pharmaceutically acceptable salt thereof.

## 13. (canceled)

14. (previously presented) A method of treating dyslipidemia comprising administering a compound of claim 1, or a pharmaceutically acceptable salt thereof, to a patient in need thereof.

# 15. (canceled)

(currently amended) A method of treating artherosclerosis comprising
 administering a compound of claim 1, or a pharmaccutically acceptable salt thereof, to a patient.

### (canceled)

 (previously presented) A method of according to claim 14 comprising lowering plasma LDL-cholesterol in a mammal.

# 19. (canceled)

20. (currently amended) A method of treating pathological sequelae due to low levels of plasma HDL-cholesterol in a mammal comprising administering a pharmaceutically effective amount of a compound of claim 1, or a pharmaceutically acceptable salt thereof, to a patient in need thereof.

# 21. (canceled)

22. (previously presented) A pharmaceutical formulation comprising a compound according to Claim 1 and at least one of: a carrier, a diluent and an excipient.

# 23-25 (canceled)

(previously presented) A method according to claim 14 comprising raising plasma
 HDL-cholesterol in a mammal.